

SPECTRAL GAPS OF THE ONE-DIMENSIONAL SCHRÖDINGER OPERATORS WITH SINGULAR PERIODIC POTENTIALS

VLADIMIR MIKHAILET'S AND VOLODYMYR MOLYBOGA

In the memory of A. Ya. Povzner.

ABSTRACT. The behaviour of the lengths of spectral gaps $\{\gamma_n(q)\}_{n=1}^{\infty}$ of the Hill-Schrödinger operators

$$S(q)u = -u'' + q(x)u, \quad u \in \text{Dom}(S(q))$$

with real-valued 1-periodic distributional potentials $q(x) \in H_{1\text{-per}}^{-1}(\mathbb{R})$ is studied. We show that they exhibit the same behaviour as the Fourier coefficients $\{\hat{q}(n)\}_{n=-\infty}^{\infty}$ of the potentials $q(x)$ with respect to the weighted sequence spaces $h^{s,\varphi}$, $s > -1$, $\varphi \in \text{SV}$. The case $q(x) \in L_{1\text{-per}}^2(\mathbb{R})$, $s \in \mathbb{Z}_+$, $\varphi \equiv 1$ corresponds to the Marchenko-Ostrovskii Theorem.

1. INTRODUCTION

The Hill-Schrödinger operators

$$S(q)u := -u'' + q(x)u, \quad u \in \text{Dom}(S(q))$$

with real-valued 1-periodic distributional potentials $q(x) \in H_{1\text{-per}}^{-1}(\mathbb{R})$ are well defined on the Hilbert space $L^2(\mathbb{R})$ in the following *equivalent* basic ways [MiMl6]:

- as form-sum operators;
- as quasi-differential operators;
- as limits of operators with smooth 1-periodic potentials in the norm resolvent sense.

The operators $S(q)$ are lower semibounded and self-adjoint on the Hilbert space $L^2(\mathbb{R})$. Their spectra are absolutely continuous and have a band and gap structure as in the classical case of $L_{1\text{-per}}^2(\mathbb{R})$ -potentials [HrMk, Krt, DjMt3, MiMl6].

The object of our investigation is the behaviour of the lengths of spectral gaps. Under the assumption

$$(1) \quad q(x) = \sum_{k \in \mathbb{Z}} \hat{q}(k) e^{ik2\pi x} \in H_{1\text{-per}}^{-1+}(\mathbb{R}, \mathbb{R}),$$

that is

$$\sum_{k \in \mathbb{Z}} (1 + 2|k|)^{2s} |\hat{q}(k)|^2 < \infty \quad \forall s > -1, \quad \text{and} \quad \text{Im } q(x) = 0,$$

we will prove many terms asymptotic estimates for the lengths $\{\gamma_n(q)\}_{n=1}^{\infty}$ and midpoints $\{\tau_n(q)\}_{n=1}^{\infty}$ of spectral gaps of the Hill-Schrödinger operators $S(q)$ (Theorem 1). These estimates enable us to establish relationship between the rate of *decreasing/increasing* of the lengths of spectral gaps and the *regularity* of the singular potentials (Theorem 2 and Theorem 3).

It is well known that if the potentials

$$q(x) = \sum_{k \in \mathbb{Z}} \hat{q}(k) e^{ik2\pi x} \in L_{1\text{-per}}^2(\mathbb{R}, \mathbb{R}),$$

2000 *Mathematics Subject Classification*. Primary 34L40; Secondary 47A10, 47A75.

Key words and phrases. Hill-Schrödinger operators, singular potentials, spectral gaps.

i.e., when

$$\sum_{k \in \mathbb{Z}} |\widehat{q}(k)|^2 < \infty, \quad \text{and} \quad \text{Im } q(x) = 0,$$

then the Hill-Schrödinger operators $S(q)$ are lower semibounded and self-adjoint operators on the Hilbert space $L^2(\mathbb{R})$ with absolutely continuous spectra which have a zone structure [DnSch2, ReSi4].

Spectra $\text{spec}(S(q))$ are defined by the location of the endpoints $\{\lambda_0(q), \lambda_n^\pm(q)\}_{n=1}^\infty$ of spectral gaps which satisfy the following inequalities:

$$-\infty < \lambda_0(q) < \lambda_1^-(q) \leq \lambda_1^+(q) < \lambda_2^-(q) \leq \lambda_2^+(q) < \dots$$

Moreover, for even/odd numbers $n \in \mathbb{Z}_+$ the endpoints of spectral gaps are eigenvalues of the periodic/semiperiodic problems on the interval $[0, 1]$:

$$\begin{aligned} S_\pm(q)u &:= -u'' + q(x)u = \lambda u, \\ \text{Dom}(S_\pm(q)) &:= \left\{ u \in H^2[0, 1] \mid u^{(j)}(0) = \pm u^{(j)}(1), j = 0, 1 \right\} \equiv H_\pm^2[0, 1]. \end{aligned}$$

Spectral bands (stability or tied zones),

$$\mathcal{B}_0(q) := [\lambda_0(q), \lambda_1^-(q)], \quad \mathcal{B}_n(q) := [\lambda_n^+(q), \lambda_{n+1}^-(q)], \quad n \in \mathbb{N},$$

are characterized as a locus of those real $\lambda \in \mathbb{R}$ for which all solutions of the equation $S(q)u = \lambda u$ are bounded. On the other hand, spectral gaps (instability or forbidden zones),

$$\mathcal{G}_0(q) := (-\infty, \lambda_0(q)), \quad \mathcal{G}_n(q) := (\lambda_n^-(q), \lambda_n^+(q)), \quad n \in \mathbb{N},$$

are a locus of those real $\lambda \in \mathbb{R}$ for which any nontrivial solution of the equation $S(q)u = \lambda u$ is unbounded.

Due to Marchenko and Ostrovskii [MrOs] the endpoints of spectral gaps of the Hill-Schrödinger operators $S(q)$ satisfy the asymptotic estimates

$$(2) \quad \lambda_n^\pm(q) = n^2\pi^2 + \widehat{q}(0) \pm |\widehat{q}(n)| + h^1(n), \quad n \rightarrow \infty.$$

As a consequence, for the lengths of spectral gaps,

$$\gamma_n(q) := \lambda_n^+ - \lambda_n^-, \quad n \in \mathbb{N},$$

the following asymptotic formulae are fulfilled,

$$(3) \quad \gamma_n(q) = 2|\widehat{q}(n)| + h^1(n), \quad n \rightarrow \infty.$$

Hochstadt [Hchs] ($\Rightarrow$) and Marchenko, Ostrovskii [MrOs], McKean, Trubowitz [McKTr] ($\Leftarrow$) proved that the potential $q(x)$ is an infinitely differentiable function if and only if the lengths of spectral gaps $\{\gamma_n(q)\}_{n=1}^\infty$ decrease faster than arbitrary power of $1/n$:

$$q(x) \in C_{1\text{-per}}^\infty(\mathbb{R}, \mathbb{R}) \Leftrightarrow \gamma_n(q) = O(n^{-k}), \quad n \rightarrow \infty \quad \forall k \in \mathbb{Z}_+.$$

Marchenko and Ostrovskii [MrOs] discovered that

$$(4) \quad q(x) \in H_{1\text{-per}}^k(\mathbb{R}, \mathbb{R}) \Leftrightarrow \{\gamma_n(q)\}_{n=1}^\infty \in h^k, \quad k \in \mathbb{Z}_+.$$

The relationship (4) was extended by Kappeler, Mityagin [KpMt] ($\Rightarrow$) and Djakov, Mityagin [DjMt1] ($\Leftarrow$) (see also the survey [DjMt2] and the references therein) on the case of symmetric, monotonic, submultiplicative and strictly subexponential weights $\Omega = \{\Omega(n)\}_{n \in \mathbb{Z}}$:

$$q(x) \in H_{1\text{-per}}^\Omega(\mathbb{R}, \mathbb{R}) \Leftrightarrow \{\gamma_n(q)\}_{n=1}^\infty \in h^\Omega.$$

Pöschel [Psch] proved the latter statement in a quite different way.

Here and throughout the remainder of the paper we use the complex Hilbert spaces $H_{1-per}^w(\mathbb{R})$ (as well as $H_{\pm}^w[0, 1]$) of 1-periodic functions and distributions defined by means of their Fourier coefficients:

$$f(x) = \sum_{k \in \mathbb{Z}} \widehat{f}(k) e^{ik2\pi x} \in H_{1-per}^w(\mathbb{R}) \Leftrightarrow \{\widehat{f}(k)\}_{k \in \mathbb{Z}} \in h^w,$$

$$h^w = \left\{ a = \{a(k)\}_{k \in \mathbb{Z}} \left| \|a\|_{h^w} = \left(\sum_{k \in \mathbb{Z}} w^2(k) |a(k)|^2 \right)^{1/2} < \infty \right. \right\}.$$

Basically we use the power weights

$$w_s = \{w_s(k)\}_{k \in \mathbb{Z}} : \quad w_s(k) := (1 + 2|k|)^s, \quad s \in \mathbb{R}.$$

The corresponding spaces we denote as

$$H_{1-per}^{w_s}(\mathbb{R}) \equiv H_{1-per}^s(\mathbb{R}), \quad H_{\pm}^{w_s}[0, 1] \equiv H_{\pm}^s[0, 1], \quad \text{and} \quad h^{w_s} \equiv h^s, \quad s \in \mathbb{R}.$$

For more details, see Appendix.

2. MAIN RESULTS

As we already remarked, under the assumption (1) the Hill-Schrödinger operators $S(q)$ are lower semibounded and self-adjoint on the Hilbert space $L^2(\mathbb{R})$. Their spectra are absolutely continuous and have a classical zone structure [HrMk, Krt, DjMt3, MiMl6, MiSb].

Using the results of the papers [KpMh, Mhr], the Isospectral Theorem 5 and [MiMl6, Theorem C] we prove uniform many terms asymptotic estimates for the lengths of spectral gaps $\{\gamma_n(q)\}_{n=1}^{\infty}$ and their midpoints $\{\tau_n(q)\}_{n=1}^{\infty}$,

$$\tau_n(q) := \frac{\lambda_n^+(q) + \lambda_n^-(q)}{2}, \quad n \in \mathbb{N}.$$

Theorem 1 ([MiMl3, Mlb2]). *Let $q(x) \in H_{1-per}^{-\alpha}(\mathbb{R}, \mathbb{R})$, $\alpha \in [0, 1)$. Then for any $\varepsilon > 0$ uniformly on the bounded sets of distributions $q(x)$ in the corresponding Sobolev spaces $H_{1-per}^{-\alpha}(\mathbb{R})$ the lengths $\{\gamma_n(q)\}_{n=1}^{\infty}$ and midpoints $\{\tau_n(q)\}_{n=1}^{\infty}$ of spectral gaps of the Hill-Schrödinger operators $S(q)$ for $n \geq n_0$ ($\|q\|_{H_{1-per}^{-\alpha}(\mathbb{R})}$) satisfy the following asymptotic formulae:*

$$(5) \quad \gamma_n(q) = 2|\widehat{q}(n)| + h^{1-2\alpha-\varepsilon}(n),$$

$$(6) \quad \tau_n(q) = n^2\pi^2 + \widehat{q}(0) + h^{1-2\alpha-\varepsilon}(n).$$

Corollary ([MiMl3, Mlb2]). *Let $q(x) \in H_{1-per}^{-\alpha}(\mathbb{R}, \mathbb{R})$ with $\alpha \in [0, 1)$. Then for any $\varepsilon > 0$ uniformly by $q(x)$ for the endpoints of spectral gaps of the Hill-Schrödinger operators $S(q)$ the following asymptotic estimates are fulfilled:*

$$\lambda_n^{\pm}(q) = n^2\pi^2 + \widehat{q}(0) \pm |\widehat{q}(n)| + h^{1-2\alpha-\varepsilon}(n).$$

Now, we can describe a bilateral relationship between the rate of decreasing/increasing of the lengths of spectral gaps $\{\gamma_n(q)\}_{n=1}^{\infty}$ and the regularity of the potentials $q(x)$ in the refined scale.

Let

$$w_{s,\varphi} = \{w_{s,\varphi}(k)\}_{k \in \mathbb{Z}} : \quad w_{s,\varphi}(k) := (1 + 2|k|)^s \varphi(|k|), \quad s \in \mathbb{R}, \quad \varphi \in \text{SV},$$

where φ is a slowly varying on $+\infty$ in a sense of Karamata function [Snt]. It means that it is a positive, measurable on $[A, \infty)$, $A > 0$ function obeying the condition

$$\lim_{t \rightarrow +\infty} \frac{\varphi(\lambda t)}{\varphi(t)} = 1, \quad \lambda > 0.$$

For example,

$$\varphi(t) = (\log t)^{r_1} (\log \log t)^{r_2} \dots (\log \dots \log t)^{r_k} \in \text{SV}, \quad \{r_1, \dots, r_k\} \subset \mathbb{R}, \quad k \in \mathbb{N}.$$

The Hörmander spaces

$$H_{1\text{-per}}^{w_{s,\varphi}}(\mathbb{R}) \equiv H_{1\text{-per}}^{s,\varphi}(\mathbb{R}) \simeq H^{s,\varphi}(\mathbb{S}), \quad \mathbb{S} := \mathbb{R}/2\pi\mathbb{Z},$$

and the weighted sequence spaces

$$h^{w_{s,\varphi}} \equiv h^{s,\varphi}$$

form the refined scales:

$$\begin{aligned} (7) \quad & H_{1\text{-per}}^{s+\varepsilon}(\mathbb{R}) \hookrightarrow H_{1\text{-per}}^{s,\varphi}(\mathbb{R}) \hookrightarrow H_{1\text{-per}}^{s-\varepsilon}(\mathbb{R}), \\ (8) \quad & h^{s+\varepsilon} \hookrightarrow h^{s,\varphi} \hookrightarrow h^{s-\varepsilon}, \quad s \in \mathbb{R}, \varepsilon > 0, \varphi \in \text{SV}, \end{aligned}$$

which in a general situation were studied by Mikhailets and Murach [MiMr].

The following statements show that the sequence $\{\gamma_n(q)\}_{n=1}^\infty$ has the same behaviour as the Fourier coefficients $\{\widehat{q}(n)\}_{n=-\infty}^\infty$ with respect to the refined scale $\{h^{s,\varphi}\}_{s \in \mathbb{R}, \varphi \in \text{SV}}$.

Theorem 2. *Let $q(x) \in H_{1\text{-per}}^{-1+}(\mathbb{R}, \mathbb{R})$. Then*

$$q(x) \in H_{1\text{-per}}^{s,\varphi}(\mathbb{R}, \mathbb{R}) \Leftrightarrow \{\gamma_n(q)\}_{n=1}^\infty \in h^{s,\varphi}, \quad s \in (-1, 0], \varphi \in \text{SV}.$$

Note that the Hörmander spaces $H_{1\text{-per}}^{s,\varphi}(\mathbb{R})$ with $\varphi \equiv 1$ coincide with the Sobolev spaces,

$$H_{1\text{-per}}^{s,1}(\mathbb{R}) \equiv H_{1\text{-per}}^s(\mathbb{R}), \quad \text{and} \quad h^{s,1} \equiv h^s, \quad s \in \mathbb{R}.$$

Corollary ([MiMl3, Mlb2]). *Let $q(x) \in H_{1\text{-per}}^{-1+}(\mathbb{R}, \mathbb{R})$, then*

$$(9) \quad q(x) \in H_{1\text{-per}}^s(\mathbb{R}, \mathbb{R}) \Leftrightarrow \{\gamma_n(q)\}_{n=1}^\infty \in h^s, \quad s \in (-1, 0].$$

Theorem 2 together with [KpMt, Theorem 1.2], and the properties (7) and (8), involve the following extension of the Marchenko-Ostrovskii Theorem (4).

Theorem 3. *Let $q(x) \in H_{1\text{-per}}^{-1+}(\mathbb{R}, \mathbb{R})$. Then*

$$q(x) \in H_{1\text{-per}}^{s,\varphi}(\mathbb{R}, \mathbb{R}) \Leftrightarrow \{\gamma_n(q)\}_{n=1}^\infty \in h^{s,\varphi}, \quad s \in (-1, \infty), \varphi \in \text{SV}.$$

In particular,

$$q(x) \in H_{1\text{-per}}^s(\mathbb{R}, \mathbb{R}) \Leftrightarrow \{\gamma_n(q)\}_{n=1}^\infty \in h^s, \quad s \in (-1, \infty).$$

Remark. In the preprint [DjMt3] the authors announced without a proof the more general statement:

$$q(x) \in H_{1\text{-per}}^{\widehat{\Omega}}(\mathbb{R}, \mathbb{R}) \Leftrightarrow \{\gamma_n(q)\}_{n=1}^\infty \in h^{\widehat{\Omega}}, \quad \widehat{\Omega} = \left\{ \frac{\Omega(n)}{1+2|n|} \right\}_{n \in \mathbb{Z}},$$

where the weights $\Omega = \{\Omega(n)\}_{n \in \mathbb{Z}}$ are supposed to be symmetric, monotonic, submultiplicative and strictly subexponential ones. This result contains the limiting case

$$q(x) \in H_{1\text{-per}}^{-1}(\mathbb{R}, \mathbb{R}) \setminus H_{1\text{-per}}^{-1+}(\mathbb{R}, \mathbb{R}).$$

An implication

$$q(x) \in H_{1\text{-per}}^{-1}(\mathbb{R}, \mathbb{R}) \Rightarrow \{\gamma_n(q)\}_{n=1}^\infty \in h^{-1}$$

was proved in the paper [Krt].

3. PROOFS

Spectra of the Hill-Schrödinger operators $S(q)$, $q(x) \in H_{1-per}^{-1}(\mathbb{R}, \mathbb{R})$ are defined by the endpoints $\{\lambda_0(q), \lambda_n^\pm(q)\}_{n=1}^\infty$ of spectral gaps. The endpoints as in the case of $L_{1-per}^2(\mathbb{R})$ -potentials satisfy the inequalities:

$$-\infty < \lambda_0(q) < \lambda_1^-(q) \leq \lambda_1^+(q) < \lambda_2^-(q) \leq \lambda_2^+(q) < \dots$$

For even/odd numbers $n \in \mathbb{Z}_+$ they are eigenvalues of the periodic/semiperiodic problems on the interval $[0, 1]$ [MiMl6, Theorem C],

$$S_\pm(q)u = \lambda u.$$

The operators

- $S_\pm u \equiv S_\pm(q)u := D_\pm^2 u + q(x)u,$
- $\bullet D_\pm^2 := -d^2/dx^2, \text{ Dom}(D_\pm^2) = H_\pm^2[0, 1];$
- $\bullet q(x) = \sum_{k \in \mathbb{Z}} \widehat{q}(k) e^{ik2\pi x} \in H_+^{-1}([0, 1], \mathbb{R});$
- $\bullet \text{Dom}(S_\pm(q)) = \{u \in H_\pm^1[0, 1] \mid D_\pm^2 u + q(x)u \in L^2(0, 1)\},$

are well defined on the Hilbert space $L^2(0, 1)$ as lower semibounded, self-adjoint form-sum operators, and they have the pure discrete spectra

$$\text{spec}(S_\pm(q)) = \{\lambda_0[S_+(q)], \lambda_{2n-1}^\pm[S_-(q)], \lambda_{2n}^\pm[S_+(q)]\}_{n=1}^\infty.$$

In the papers [MiMl3, Mlb2, MiMl4, MiMl5] the authors meticulously investigated the more general periodic/semiperiodic form-sum operators

$$S_{m,\pm}(V) := D_\pm^{2m} + V(x), \quad V(x) \in H_+^{-m}[0, 1], \quad m \in \mathbb{N},$$

on the Hilbert space $L^2(0, 1)$.

So, we need to find precise asymptotic estimates for the operators $S_\pm(q)$ eigenvalues. It is quite difficult problem as the form-sum operators $S_\pm(q)$ are not convenient for investigation. We also cannot apply approach developed by Savchuk and Shkalikov (see the survey [SvSh] and the references therein) considering the operators $S_\pm(q)$ as quasi-differential ones as the periodic/semiperiodic boundary conditions are not strongly regular by Birkhoff. Therefore we propose an alternative approach which is based on isospectral transformation of the problem.

Kappeler and Möhr [KpMh, Mhr] investigated the second order differential operators $L_\pm(q)$, $q(x) \in H_+^{-1}([0, 1], \mathbb{R})$ (in general, with complex-valued potentials) defined on the *negative* Sobolev spaces $H_\pm^{-1}[0, 1]$,

$$L_\pm \equiv L_\pm(q) := D_\pm^2 + q(x), \quad \text{Dom}(L_\pm(q)) = H_\pm^1[0, 1].$$

They established that the operators $L_\pm(q)$ with $q(x) \in H_+^{-\alpha}([0, 1], \mathbb{R})$, $\alpha \in [0, 1)$ have the real-valued discrete spectra

$$\text{spec}(L_\pm(q)) = \{\lambda_0[L_+(q)], \lambda_{2n-1}^\pm[L_-(q)], \lambda_{2n}^\pm[L_+(q)]\}_{n=1}^\infty$$

such that

$$|\lambda_n^\pm[L_\pm(q)] - n^2\pi^2 - \widehat{q}(0)| \leq Cn^\alpha, \quad n \geq n_0 \left(\|q\|_{H_+^{-\alpha}[0, 1]} \right).$$

More precisely, for the values

$$\begin{aligned} \gamma_n[L_\pm(q)] &:= \lambda_n^+[L_\pm(q)] - \lambda_n^-[L_\pm(q)], \quad n \in \mathbb{N}, \\ \tau_n[L_\pm(q)] &:= \frac{\lambda_n^+[L_\pm(q)] + \lambda_n^-[L_\pm(q)]}{2}, \quad n \in \mathbb{N} \end{aligned}$$

they proved the following result.

Proposition 4 (Kappeler, Möhr [KpMh, Mhr]). *Let $q(x) \in H_+^{-\alpha}([0, 1], \mathbb{R})$, and $\alpha \in [0, 1]$. Then for any $\varepsilon > 0$ uniformly on the bounded sets of distributions $q(x)$ in the Sobolev spaces $H_+^{-\alpha}[0, 1]$ for the operators $L_{\pm}(q)$ values $\{\gamma_n[L_{\pm}(q)]\}_{n=1}^{\infty}$ and $\{\tau_n[L_{\pm}(q)]\}_{n=1}^{\infty}$ for $n \geq n_0 \left(\|q\|_{H_+^{-\alpha}[0, 1]} \right)$ the following asymptotic estimates are fulfilled:*

$$\begin{aligned} i) \quad & \left\{ \min_{\pm} \left| \gamma_n[L_{\pm}(q)] \pm 2\sqrt{(\widehat{q} + \omega)(-n)(\widehat{q} + \omega)(n)} \right| \right\}_{n \in \mathbb{N}} \in h^{1-2\alpha-\varepsilon}, \\ ii) \quad & \tau_n[L_{\pm}(q)] = n^2\pi^2 + \widehat{q}(0) + h^{1-2\alpha-\varepsilon}(n), \end{aligned}$$

where the convolution

$$\{\omega(n)\}_{n \in \mathbb{Z}} \equiv \left\{ \frac{1}{\pi^2} \sum_{k \in \mathbb{Z} \setminus \{\pm n\}} \frac{\widehat{q}(n-k)\widehat{q}(n+k)}{n^2 - k^2} \right\}_{n \in \mathbb{Z}} \in \begin{cases} h^{1-\alpha}, & \alpha \in [0, 1/2), \\ h^{3/2-2\alpha-\delta}, & \alpha \in [1/2, 1) \end{cases}$$

with any $\delta > 0$ (see the Convolution Lemma [KpMh, Mhr]).

Remark. In the papers [Mlb1, MiMl1, MiMl2, Mlb2] more general operators

$$L_{m,\pm}(V) := D_{\pm}^{2m} + V(x), \quad V(x) \in H_+^{-m}[0, 1], \quad m \in \mathbb{N}$$

on the spaces $H_{\pm}^{-m}[0, 1]$ were studied. In particular, the analogue of Proposition 4 was proved.

The following statement is an essential point of our approach.

Theorem 5 (Isospectral Theorem [MiMl3, Mlb2]). *The operators $S_{\pm}(q)$ and $L_{\pm}(q)$ are isospectral ones:*

$$\text{spec}(S_{\pm}(q)) = \text{spec}(L_{\pm}(q)).$$

Proof. The injections

$$\text{spec}(S_{\pm}(q)) \subset \text{spec}(L_{\pm}(q))$$

are obvious since

$$S_{\pm}(q) \subset L_{\pm}(q).$$

Let prove the inverse injections

$$\text{spec}(L_{\pm}(q)) \subset \text{spec}(S_{\pm}(q)).$$

Let $\lambda \in \text{spec}(L_{\pm}(q))$, and f be a correspondent eigenvector or rootvector. Therefore

$$(L_{\pm}(q) - \lambda Id)f = g, \quad f, g \in \text{Dom}(L_{\pm}(q)) = H_{\pm}^1[0, 1],$$

where f is an eigenfunction if $g = 0$, and a rootvector if $g \neq 0$.

So, we have got

$$L_{\pm}(q)f = \lambda Idf + g \in H_{\pm}^1[0, 1],$$

i.e.,

$$L_{\pm}(q)f = D_{\pm}^2 f + q(x)f \in L^2(0, 1).$$

Thus we have proved that $f \in \text{Dom}(S_{\pm}(q))$. In the case when f is a rootvector ($g \neq 0$) in a similar fashion we show that $g \in \text{Dom}(S_{\pm}(q))$ also. Continuing this process until necessary (draw attention that it is finite as the eigenvalue λ has a finite algebraic multiplicity) we obtain that all correspondent to λ eigenvectors and rootvectors belong to the operators $S_{\pm}(q)$ domains $\text{Dom}(S_{\pm}(q))$. Consequently we can conclude that

$$\lambda \in \text{spec}(S_{\pm}(q)),$$

and the required injections

$$\text{spec}(L_{\pm}(q)) \subset \text{spec}(S_{\pm}(q))$$

have been proved.

The proof is complete. $\square$

Now, Theorem 1 follows from Proposition 4, the Isospectral Theorem 5 and [MiMl6, Theorem C], since

$$\begin{aligned}\widehat{q}(n) &= \overline{\widehat{q}(-n)}, \quad n \in \mathbb{Z}, \\ \omega(n) &= \overline{\omega(-n)}, \quad n \in \mathbb{Z},\end{aligned}$$

and as a consequence

$$\min_{\pm} \left| \gamma_n(q) \pm 2\sqrt{(\widehat{q} + \omega)(-n)(\widehat{q} + \omega)(n)} \right| = |\gamma_n(q) - 2|(\widehat{q} + \omega)(n)||.$$

The proof of Theorem 1 is complete.

To prove Theorem 2 we firstly prove its Corollary. The formula (9) follows from [KpMh, Corollary 0.2 (2.6)], the Isospectral Theorem 5 and [MiMl6, Theorem C]. Also it can be proved directly as well as [KpMh, Corollary 0.2 (2.6)] using the estimates (5).

Further, to prove Theorem 2 it is sufficient to apply the asymptotic estimates (5), the properties (7) and (8) of the refined scales, and the formula (9):

$$\begin{aligned}\bullet \quad q &\in H_{1-per}^{s,\varphi}(\mathbb{R}, \mathbb{R}) \xrightarrow{(7)} q \in H_{1-per}^{s-\delta}(\mathbb{R}, \mathbb{R}), \delta > 0 \xrightarrow{(5)} \gamma_n = 2|\widehat{q}(n)| + h^{1+2(s-\delta)-\varepsilon}(n) \\ &\xrightarrow{(8)} \gamma_n = 2|\widehat{q}(n)| + h^{s,\varphi}(n) \implies \{\gamma_n(q)\}_{n=1}^\infty \in h^{s,\varphi}; \\ \bullet \quad \{\gamma_n(q)\}_{n=1}^\infty &\in h^{s,\varphi} \xrightarrow{(8)} \{\gamma_n\}_{n=1}^\infty \in h^{s-\delta}, \delta > 0 \xrightarrow{(9)} q \in H_{1-per}^{s-\delta}(\mathbb{R}, \mathbb{R}) \\ &\xrightarrow{(5)} \gamma_n = 2|\widehat{q}(n)| + h^{1+2(s-\delta)-\varepsilon}(n) \xrightarrow{(8)} \gamma_n = 2|\widehat{q}(n)| + h^{s,\varphi}(n) \\ &\implies \{\widehat{q}(n)\}_{n \in \mathbb{Z}} \in h^{s,\varphi}(n).\end{aligned}$$

Remark that due to arbitrary choice of $\delta > 0$ and $\varepsilon > 0$ we may choose them such that

$$1 + s - 2\delta - \varepsilon > 0.$$

The proof of Theorem 2 is complete. Now, we are ready to prove Theorem 3.

At first, note that from [KpMt, Theorem 1.2] we get the following asymptotic formulae for the lengths of spectral gaps:

$$(10) \quad \gamma_n(q) = 2|\widehat{q}(n)| + h^{1+s}(n) \quad \text{as } q(x) \in H_{1-per}^s(\mathbb{R}, \mathbb{R}), \quad s \in [0, \infty),$$

which for the integer numbers $s \in \mathbb{Z}_+$ were proved by Marchenko and Ostrovskii [MrOs].

Using (9), (10) and (4) it is easy to prove the relationship

$$(11) \quad q(x) \in H_{1-per}^s(\mathbb{R}, \mathbb{R}) \Leftrightarrow \{\gamma_n(q)\}_{n=1}^\infty \in h^s, \quad s \in (-1, \infty).$$

Sufficiency of Theorem 3. Let $q(x) \in H_{1-per}^{s,\varphi}(\mathbb{R}, \mathbb{R})$. If $s \in (-1, 0]$ then due to Theorem 2 we obtain that $\{\gamma_n(q)\}_{n=1}^\infty \in h^{s,\varphi}$. If $s > 0$ then

$$\begin{aligned}q(x) &\in H_{1-per}^{s,\varphi}(\mathbb{R}, \mathbb{R}) \xrightarrow{(7)} H_{1-per}^{s-\delta}(\mathbb{R}, \mathbb{R}), \delta > 0 \xrightarrow{(10)} \gamma_n(q) = 2|\widehat{q}(n)| + h^{1+s-\delta}(n) \\ &\xrightarrow{(8)} \gamma_n(q) = 2|\widehat{q}(n)| + h^{s,\varphi}(n) \implies \{\gamma_n(q)\}_{n=1}^\infty \in h^{s,\varphi}.\end{aligned}$$

Sufficiency is proved.

Necessity of Theorem 3. Let suppose that $\{\gamma_n(q)\}_{n=1}^\infty \in h^{s,\varphi}$. If $s \in (-1, 0]$ then from Theorem 2 we have that $q(x) \in H_{1-per}^{s,\varphi}(\mathbb{R}, \mathbb{R})$. If $s > 0$ then

$$\begin{aligned}\{\gamma_n(q)\}_{n=1}^\infty &\in h^{s,\varphi} \xrightarrow{(8)} h^{s-\delta}, \delta > 0 \xrightarrow{(11)} q(x) \in H_{1-per}^{s-\delta}(\mathbb{R}, \mathbb{R}) \xrightarrow{(10)} \gamma_n(q) = 2|\widehat{q}(n)| + h^{1+s-\delta}(n) \\ &\xrightarrow{(8)} \gamma_n(q) = 2|\widehat{q}(n)| + h^{s,\varphi}(n) \implies q(x) \in H_{1-per}^{s,\varphi}(\mathbb{R}, \mathbb{R}).\end{aligned}$$

Necessity is proved.

The proof of Theorem 3 is complete.

4. CONCLUDING REMARKS

In fact we can prove the following result: if $q(x) \in H_{1-per}^{-1+}(\mathbb{R}, \mathbb{R})$ and

$$\begin{aligned} (1 + 2|k|)^s &\ll w(k) \ll (1 + 2|k|)^{1+2s}, \quad s \in (-1, 0], \\ (1 + 2|k|)^s &\ll w(k) \ll (1 + 2|k|)^{1+s}, \quad s \in [0, \infty), \end{aligned}$$

then

$$q(x) \in H_{1-per}^w(\mathbb{R}, \mathbb{R}) \Leftrightarrow \{\gamma_n(q)\}_{n=1}^\infty \in h^w.$$

This result is not covered by the theorems of the preprint [DjMt3], because it does not require the weight function to be monotonic and submultiplicative.

APPENDIX

The complex Sobolev spaces $H_{1-per}^s(\mathbb{R})$, $s \in \mathbb{R}$ of 1-periodic functions and distributions over the real axis $\mathbb{R}$ are defined by means of their Fourier coefficients,

$$\begin{aligned} H_{1-per}^s(\mathbb{R}) &:= \left\{ f = \sum_{k \in \mathbb{Z}} \widehat{f}(k) e^{ik2\pi x} \mid \|f\|_{H_{1-per}^s(\mathbb{R})} < \infty \right\}, \\ \|f\|_{H_{1-per}^s(\mathbb{R})} &:= \left(\sum_{k \in \mathbb{Z}} \langle 2k \rangle^{2s} |\widehat{f}(k)|^2 \right)^{1/2}, \quad \langle k \rangle := 1 + |k|, \\ \widehat{f}(k) &:= \langle f, e^{ik2\pi x} \rangle_{L_{1-per}^2(\mathbb{R})}, \quad k \in \mathbb{Z}. \end{aligned}$$

By $\langle \cdot, \cdot \rangle_{L_{1-per}^2(\mathbb{R})}$ we denote a sesqui-linear form pairing the dual spaces $H_{1-per}^s(\mathbb{R})$ and $H_{1-per}^{-s}(\mathbb{R})$ with respect to $L_{1-per}^2(\mathbb{R})$, which (the sequi-linear form $\langle \cdot, \cdot \rangle_{L_{1-per}^2(\mathbb{R})}$) is an extension by continuity of the $L_{1-per}^2(\mathbb{R})$ -inner product [Brz, GrGr]:

$$\langle f, g \rangle_{L_{1-per}^2(\mathbb{R})} := \int_0^1 f(x) \overline{g(x)} dx = \sum_{k \in \mathbb{Z}} \widehat{f}(k) \overline{\widehat{g}(k)} \quad \forall f, g \in L_{1-per}^2(\mathbb{R}).$$

It is useful to notice that

$$H_{1-per}^0(\mathbb{R}) \equiv L_{1-per}^2(\mathbb{R}).$$

By $H_{1-per}^{s+}(\mathbb{R})$, $s \in \mathbb{R}$ we denote an inductive limit of the Sobolev spaces $H_{1-per}^t(\mathbb{R})$ with $t > s$,

$$H_{1-per}^{s+}(\mathbb{R}) := \bigcup_{\varepsilon > 0} H_{1-per}^{s+\varepsilon}(\mathbb{R}).$$

It is a topological space with an inductive topology.

In a similar fashion the Sobolev spaces $H_{\pm}^s[0, 1]$, $s \in \mathbb{R}$ of 1-periodic/1-semiperiodic functions and distributions over the interval $[0, 1]$ are defined:

$$\begin{aligned} H_{\pm}^s[0, 1] &:= \left\{ f = \sum_{k \in \Gamma_{\pm}} \widehat{f}\left(\frac{k}{2}\right) e^{ik\pi x} \mid \|f\|_{H_{\pm}^s[0, 1]} < \infty \right\}, \\ \|f\|_{H_{\pm}^s[0, 1]} &:= \left(\sum_{k \in \Gamma_{\pm}} \langle k \rangle^{2s} \left| \widehat{f}\left(\frac{k}{2}\right) \right|^2 \right)^{1/2}, \quad \langle k \rangle = 1 + |k|, \\ \widehat{f}\left(\frac{k}{2}\right) &:= \langle f(x), e^{ik\pi x} \rangle_{\pm}, \quad k \in \Gamma_{\pm}. \end{aligned}$$

Here

$$\begin{aligned}\Gamma_+ &\equiv 2\mathbb{Z} := \{k \in \mathbb{Z} \mid k \equiv 0 \pmod{2}\}, \\ \Gamma_- &\equiv 2\mathbb{Z} + 1 := \{k \in \mathbb{Z} \mid k \equiv 1 \pmod{2}\},\end{aligned}$$

and $\langle \cdot, \cdot \rangle_{\pm}$ are sesqui-linear forms pairing the dual spaces $H_{\pm}^s[0, 1]$ and $H_{\pm}^{-s}[0, 1]$ with respect to $L^2(0, 1)$, which (the sesqui-linear forms $\langle \cdot, \cdot \rangle_{\pm}$) are extensions by continuity of the $L^2(0, 1)$ -inner product [Brz, GrGr]:

$$\langle f, g \rangle_{\pm} := \int_0^1 f(x) \overline{g(x)} dx = \sum_{k \in \Gamma_{\pm}} \widehat{f}\left(\frac{k}{2}\right) \overline{\widehat{g}\left(\frac{k}{2}\right)} \quad \forall f, g \in L^2(0, 1).$$

It is obvious that

$$H_+^0[0, 1] \equiv H^0[0, 1] \equiv L^2(0, 1).$$

We say that 1-periodic function or distribution $f(x)$ is *real-valued* if $\text{Im } f(x) = 0$. Let us remind that

$$\text{Re } f(x) := \frac{1}{2}(f(x) + \overline{f(x)}), \quad \text{Im } f(x) := \frac{1}{2i}(f(x) - \overline{f(x)}),$$

(see, for an example, [Vld]). In terms of the Fourier coefficients we have

$$\text{Im } f(x) = 0 \Leftrightarrow \widehat{f}(k) = \overline{\widehat{f}(-k)}, \quad k \in \mathbb{Z}.$$

Set

$$\begin{aligned}H_{1\text{-per}}^s(\mathbb{R}, \mathbb{R}) &:= \{f(x) \in H_{1\text{-per}}^s(\mathbb{R}) \mid \text{Im } f(x) = 0\}, \\ H_{1\text{-per}}^{s+}(\mathbb{R}, \mathbb{R}) &:= \{f(x) \in H_{1\text{-per}}^{s+}(\mathbb{R}) \mid \text{Im } f(x) = 0\}, \\ H_{\pm}^s([0, 1], \mathbb{R}) &:= \{f(x) \in H_{\pm}^s[0, 1] \mid \text{Im } f(x) = 0\}.\end{aligned}$$

Also we will need the Hilbert sequence spaces

$$h^s \equiv h^s(\mathbb{Z}, \mathbb{C}), \quad s \in \mathbb{R}$$

of (two-sided) weighted sequences,

$$h^s := \left\{ a = \{a(k)\}_{k \in \mathbb{Z}} \mid \|a\|_{h^s} := \left(\sum_{k \in \mathbb{Z}} \langle k \rangle^{2s} |a(k)|^2 \right)^{1/2} < \infty \right\}, \quad \langle k \rangle = 1 + |k|.$$

Note that

$$h^0 \equiv l^2(\mathbb{Z}, \mathbb{C}),$$

and

$$a = \{a(k)\}_{k \in \mathbb{Z}} \in h^s, \quad s \in \mathbb{R} \quad \Rightarrow \quad a(k) = o(|k|^{-s}), \quad k \rightarrow \pm\infty.$$

REFERENCES

- [Brz] Yu. M. Berezanskii, *Expansions in Eigenfunctions of Self-Adjoint Operators*, Transl. Math. Monographs, vol. 17, Amer. Math. Soc., Providence, R. I, 1968. (Russian edition: Naukova Dumka, Kiev, 1965)
- [DjMt1] P. Djakov, B. Mityagin, *Spectral triangles of Schrödinger operators with complex potentials*, Selecta Math. (N.S.) **9** (2003), no. 4, 495–528.
- [DjMt2] P. Djakov, B. Mityagin, *Instability zones of one-dimensional periodic Schrödinger and Dirac operators*, Uspekhi Mat. Nauk **61** (2006), no. 4, 77–182. (Russian)
- [DjMt3] P. Djakov, B. Mityagin, *Fourier method for one dimensional Schrödinger operators with singular periodic potentials*, arXiv:math.SP/0710.0237v1 1 Oct 2007, 1–39.
- [DnSch2] N. Dunford, J. T. Schwartz, *Linear Operators*, Part II: *Spectral Theory. Self-Adjoint Operators in Hilbert Space*, Interscience, New York, London, 1963. (Russian edition: Mir, Moscow, 1966)
- [Edv1] R. Edwards, *Fourier Series: A Modern Introduction*. Vol. I, Holt, Rinehard and Winston, Inc., New York-Montreal, Que.-London, 1967. (Russian edition: Mir, Moscow, 1985)
- [Edv2] R. Edwards, *Fourier Series: A Modern Introduction*. Vol. II, Holt, Rinehard and Winston, Inc., New York-Montreal, Que.-London, 1967. (Russian edition: Mir, Moscow, 1985)

- [GrGr] V. I. Gorbachuk, M. L. Gorbachuk, *Boundary Value Problems for Operator Differential Equations*, Kluwer, Dordrecht, etc., 1991. (Russian edition: Naukova Dumka, Kiev, 1984)
- [HrMk] R. Hryniv, Ya. Mykytyuk, *1-D Schrödinger operators with periodic singular potentials*, Methods Funct. Anal. Topology **7** (2001), no. 4, 31–42.
- [Hchs] H. Hochstadt, *Estimates of the stability intervals for Hill's equation*, Proc. Amer. Math. Soc. **14** (1963), 930–932.
- [KpMt] T. Kappeler, B. Mityagin, *Estimates for periodic and Dirichlet eigenvalues of the Schrödinger operator*, SIAM J. Math. Anal. **33** (2001), no. 1, 113–152.
- [KpMh] T. Kappeler, C. Möhr, *Estimates for periodic and Dirichlet eigenvalues of the Schrödinger operator with singular potentials*, J. Funct. Anal. **186** (2001), 62–91.
- [Krt] E. Korotyaev, *Characterization of the spectrum of Schrödinger operators with periodic distributions*, Int. Math. Res. Not. **37** (2003), 2019–2031.
- [MrOs] V. Marchenko, I. Ostrovskii, *A characterization of the spectrum of Hill's operator*, Matem. Sbornik **97** (1975), no. 4, 540–606. (Russian)
- [McKTr] H. McKean, E. Trubowitz, *Hill's operators and hyperelliptic function theory in the presence of infinitely many branch points*, Comm. Pure Appl. Math. **29** (1976), no. 2, 143–226.
- [MiM1] V. Mikhailets, V. Molyboga, *Singular eigenvalue problems on the circle*, Methods Funct. Anal. Topology **10** (2004), no. 3, 44–53.
- [MiM2] V. Mikhailets, V. Molyboga, *Uniform estimates for the semi-periodic eigenvalues of the singular differential operators*, Methods Funct. Anal. Topology **10** (2004), no. 4, 30–57.
- [MiM3] V. Mikhailets, V. Molyboga, *The spectral problems over the periodic classes of distributions*, Preprint, Institute of Mathematics of NAS of Ukraine, Kyiv, 2004, 46 p. (Ukrainian)
- [MiM4] V. Mikhailets, V. Molyboga, *The perturbation of periodic and semiperiodic operators by Schwartz distributions*, Reports of NAS of Ukraine **7** (2006), 26 – 31. (Russian)
- [MiM5] V. Mikhailets, V. Molyboga, *Singularly perturbed periodic and semiperiodic differential operators*, Ukrainian Math. J. **59** (2007), no. 6, 785–797.
- [MiM6] V. Mikhailets, V. Molyboga, *One-dimensional Schrödinger operators with singular periodic potentials*, Methods Funct. Anal. Topology **14** (2008), no. 2, 184–200.
- [MiMr] V. Mikhailets, A. Murach, *Interpolation with a function parameter and refined scale of spaces*, Methods Funct. Anal. Topology **14** (2008), no. 1, 81–100.
- [MiSb] V. Mikhailets, A. Sobolev, *Common eigenvalue problem and periodic Schrödinger operators*, J. Funct. Anal. **165** (1999), no. 1, 150–172.
- [Mlb1] V. Molyboga, *Estimates for periodic eigenvalues of the differential operator $(-1)^m d^{2m}/dx^{2m} + V$ with V – distribution*, Methods Funct. Anal. Topology **9** (2003), no. 2, 163–178.
- [Mlb2] V. Molyboga, *Spectra of periodic problems with generalized functions*, Candidate Dissertation, Kyiv, 2005, 167 p. (Ukrainian)
- [Mhr] C. Möhr, *Schrödinger Operators with Singular Potentials on the Circle: Spectral Analysis and Applications*, Thesis at the University of Zürich, 2001, 134 p.
- [Psch] J. Pöschel, *Hill's potentials in weighted Sobolev spaces and their spectral gaps*, Preprint, Stuttgarter Mathematische Berichte, Stuttgart, 2004, 28 p.
- [ReSi4] M. Reed, B. Simon, *Methods of Modern Mathematical Physics: Vols 1-4*, Academic Press, New York, etc., 1972–1978, V. 4: *Analysis of Operators*, 1972. (Russian edition: Mir, Moscow, 1982)
- [SvSh] A. Savchuk, A. Shkalikov, *Sturm-Liouville operators with distribution potentials*, Tr. Mosk. Mat. Obs. **64** (2003), 159–212. (Russian) Engl. transl. in Trans. Moscow Math. Soc. (2003), 143–192.
- [Snt] E. Seneta, *Regularly Varying Functions*, Lecture Notes in Math., vol. 508, Springer-Verlag, Berlin, 1976. (Russian edition: Nauka, Moscow, 1985)
- [Vld] V. S. Vladimirov, *Generalized Functions in Mathematical Physics*, Nauka, Moscow, 1976, 280 p. (Russian)

INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCE OF UKRAINE, TERESHCHENKIVSKA STR.,
3, KYIV-4, UKRAINE, 01601
E-mail address: mikhailets@imath.kiev.ua

INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCE OF UKRAINE, TERESHCHENKIVSKA STR.,
3, KYIV-4, UKRAINE, 01601
E-mail address: molyboga@imath.kiev.ua